

Physics 224

The Interstellar Medium

Lecture #8: HII Regions



Credit: NASA,ESA, M. Robberto (Space Telescope Science Institute/ESA)
and the Hubble Space Telescope Orion Treasury Project Team

Outline

- Part I: HII Regions
- Part II: Collisional Excitation
- Part III: Nebular Diagnostics
- Part IV: Heating & Cooling in HII Regions

HII Regions

Stromgren's insight:

HII regions surrounding massive, young stars are regions where H is ~fully ionized with a sharp boundary.

We heard on Friday about the full calculation that shows this is the case.

We can use this to estimate HII region properties as follows...

HII Regions

Stromgren's insight:

HII regions surrounding massive, young stars are regions where H is ~fully ionized with a sharp boundary.

ionizing photon
production rate

$\approx$

recombination rate

$$Q_0 = \frac{4\pi}{3} R_{SO}^3 \alpha_B n(H^+) n_e$$

ionizing
photons per sec

volume of
Stromgren sphere

recomb rate
per volume
Case B!

HII Regions

$$R_{S0} = \left(\frac{3Q_0}{4\pi n_H^2 \alpha_B} \right)^{1/3} = 9.77 \times 10^{18} Q_{0,49}^{1/3} n_2^{-2/3} T_4^{0.28} \text{cm}$$

where:

$$Q_{0,49} = Q_0 / 10^{49} \text{s}^{-1}$$

$$n_2 = n_H / 10^2 \text{cm}^{-3}$$

$$T_4 = T / 10^4 \text{K}$$

At n_2 , T_4 , and $Q_{0,49}$

$$R_{S0} \sim 3 \text{pc}$$

Decreases in size when n increases.
Increases when Q_0 increases.

HII Regions

Stromgren's insight:

HII regions surrounding massive, young stars are regions where H is ~fully ionized with a sharp boundary.

Transition from ionized to neutral will be approximately the mean free path of ionizing photons in HI.

$$l_{\text{mfp}} = \frac{1}{n(H^0) \sigma_{pi}} = 3.39 \times 10^{17} \left(\frac{n(H^0)}{1 \text{ cm}^{-3}} \right)^{-1} \text{ cm}$$

here: mfp for 18 eV photon

HII Regions

Calculation from Stromgren (& ch 15.3 in Draine)
of ionization fraction as a function of radius

$$\frac{x^2}{1-x} \approx \frac{1-y^3}{3y^2} \tau_S$$

$x = n_e/n_H$
 $y = r/R_{S0}$
 $\tau_S = n_H \sigma_{pi} R_{S0}$

Can calculate "typical" value from radius where 1/2 of mass is enclosed

$$(1 - x_m) = 1.1 \times 10^{-3} Q_{0,49}^{-1/3} n_2^{-1/3}$$

HII Regions

Timescale for ionization is short:


H to ionize

$$\tau_{\text{ioniz}} \equiv \frac{(4/3)\pi R_{S0}^3 n_H}{Q_0} = \frac{1}{\alpha_B n_H} = \frac{1.22 \times 10^3 \text{yr}}{n_2}$$

ionizing photons per sec

Ionization equilibrium happens quickly after star turns on.


HII Regions




HII regions have more than just H in them, ionization structure in other elements depends on stellar spectrum and density.

HII Regions

Next abundant element: He
Ionization potential 24.59 eV




"Cool" < 40,000 K star




HII Regions

Next abundant element: He
Ionization potential 24.59 eV




"Hot" 40-100,000 K star



HII Regions

Next abundant element: He
Ionization potential 24.59 eV



For stars or ionizing sources
with enough photons at $E > 54.4$ eV
get He^{++} zone

HII Regions

Photoionization Modeling:
coupling of ionization state, stellar spectrum,
density, temperature, etc for multiple species

Cloudy & Associates

Photoionization Simulations for the Discriminating Astrophysicist Since 1978

Part II: Collisional Excitation


Collisional Excitation

is important because:

- 1) it can put electrons in excited states that radiatively decay and remove energy from the gas
- 2) radiative transitions fed by collisional excitation give us very useful diagnostics of gas conditions

Collisional Excitation

Two Level Atom



Assume no background
radiation field
(i.e. ignore stimulated emission)


$$\frac{dn_1}{dt} = (\text{rate of collisions from 0 to 1}) -$$
$$(\text{rate of collisions from 1 to 0}) -$$
$$(\text{spontaneous emission from 1 to 0})$$

$$\frac{dn_1}{dt} = n_c n_0 k_{01} -$$
$$n_c n_1 k_{10} -$$
$$n_1 A_{10}$$

*per volume

Collisional Excitation

Two Level Atom



In steady state:
 $dn_1/dt = 0$

$$\frac{n_1}{n_0} = \frac{n_c k_{01}}{n_c k_{10} + A_{10}}$$

from detailed balance: $k_{01} = \frac{g_1}{g_0} k_{10} e^{-E_{10}/kT_{gas}}$

Collisional Excitation

$$\frac{n_1}{n_0} = \left(\frac{1}{1 + A_{10}/(n_c k_{10})} \right) \frac{g_1}{g_0} e^{-E_{10}/kT_{\text{gas}}}$$

define “critical density”

ratio of collisional to spontaneous rates
that depopulate level 1

$$n_{\text{crit}} = \frac{A_{10}}{k_{10}}$$

$$\frac{n_1}{n_0} = \left(\frac{1}{1 + n_{\text{crit}}/n_c} \right) \frac{g_1}{g_0} e^{-E_{10}/kT_{\text{gas}}}$$

Collisional Excitation

$$\frac{n_1}{n_0} = \left(\frac{1}{1 + n_{\text{crit}}/n_c} \right) \frac{g_1}{g_0} e^{-E_{10}/kT_{\text{gas}}}$$

When $n_c \gg n_{\text{crit}}$, level populations are set by the gas temperature and degeneracy - "thermalized"

When $n_c \ll n_{\text{crit}}$, factor in parenthesis goes to n_c/n_{crit} , population in level n_1 is "sub-thermal"

Collisional Excitation

General formulation takes into account stimulated emission and absorption too...

$$\frac{n_1}{n_0} = \frac{n_c k_{01} + \langle n_\gamma \rangle (g_1/g_0) A_{10}}{n_c k_{10} + (1 + \langle n_\gamma \rangle) A_{10}},$$

general definition of n_{crit} :

$$n_{\text{crit}} = \frac{(1 + \langle n_\gamma \rangle) A_{10}}{k_{10}}$$

where:

$$\langle n_\gamma \rangle = \frac{c^3}{8\pi h \nu^3} u_\nu$$

is the photon occupation number

Collisional Excitation

Useful to rewrite this with brightness temperature:

$$\langle n_\gamma \rangle = \frac{1}{e^{h\nu/kT_B} - 1}$$

$$\frac{n_1}{n_0} = \left(\frac{1}{1 + n_{\text{crit}}/n_c} \right) \frac{g_1}{g_0} e^{-E_{10}/kT_{\text{gas}}} + \left(\frac{1}{1 + n_c/n_{\text{crit}}} \right) \frac{g_1}{g_0} e^{-h\nu/kT_B}$$

Ratio of n_c/n_{crit} determines if level populations track gas temperature or radiation field temperature!

Critical Density

Multi-level atoms

$$n_{\text{crit},u}(c) \equiv \frac{\sum_{l < u} [1 + \langle n_{\gamma} \rangle_{ul}] A_{ul}}{\sum_{l < u} k_{ul}(c)}$$

ratio of total radiative and collisional
depopulation rates to lower levels

note: only good in cases where gas is optically
thin to radiation from $u \rightarrow l$ transition

Part III: Nebular Diagnostics

Nebular Diagnostics

Collisionally excited lines from ionized gas that give us diagnostics for density, temperature, etc.

Two types:

- 1) temperature sensitive
- 2) density sensitive

Nebular Diagnostics

Element	H II and He I zone ^b		H II and He II zone ^c	
	Ion	$h\nu$ (eV) ^d	Ion	$h\nu$ (eV) ^d
H	H II	13.60	H II	13.60
He	He I	0	He II	24.59
C	C II	11.26	C III ^e	24.38
			C IV	47.88
N	N II	14.53	N III	29.60
			N IV	47.45
O	O II	13.62	O III	35.12
Ne	Ne II	21.56	Ne III	40.96
Na	(Na II) ^f	5.14	(Na II) ^f	5.14
			Na III	47.29
Mg	Mg II	7.65	(Mg III) ^f	15.04
	(Mg III) ^f	15.04		
Al	Al III	18.83	(Al IV) ^f	28.45
Si	Si III	16.35	Si IV	33.49
			(Si V) ^f	45.14
S	S II	10.36	S III	23.33
	S III	23.33	S IV	34.83
Ar	Ar II	15.76	Ar III	27.63
			Ar IV	40.74
Ca	Ca III	11.87	Ca IV	50.91
Fe	Fe III	16.16	Fe IV	30.65
Ni	Ni III	18.17	Ni IV	35.17

^a Limited to elements X with $N_X/N_H > 10^{-6}$.

^b Ions that can be created by radiation with $13.60 < h\nu < 24.59$ eV.

^c Ions that can be created by radiation with $24.59 < h\nu < 54.42$ eV.

^d Photon energy required to create ion.

^e Ionization potential is just below 24.59 eV.

^f Closed shell, with no excited states below 13.6 eV.

First good to note which atoms and ions will be abundant in HII regions.

Temperature Sensitive Line Ratios



What we want:

two levels that can both be collisionally excited at typical HII region temperatures ($\sim 10^4$ K) but which have different enough energies that the ratio of populations depends on temperature of the gas

Requires two energy levels with $E/k < 70,000$ K

Temperature Sensitive Line Ratios

best candidates: np^2 & np^4



Temperature Sensitive Line Ratios

Ground configuration	Terms (in order of increasing energy)	Examples
$\dots ns^1$	$^2S_{1/2}$	H I, He II, C IV, N V, O VI
$\dots ns^2$	1S_0	He I, C III, N IV, O V
$\dots np^1$	$^2P_{1/2,3/2}^o$	C II, N III, O IV
$\dots np^2$	$^3P_{0,1,2}, ^1D_2, ^1S_0$	C I, N II, O III, Ne V, S III
$\dots np^3$	$^4S_{3/2}^o, ^2D_{3/2,5/2}^o, ^2P_{1/2,3/2}^o$	N I, O II, Ne IV, S II, Ar IV
$\dots np^4$	$^3P_{2,1,0}, ^1D_2, ^1S_0$	O I, Ne III, Mg V, Ar III
$\dots np^5$	$^2P_{3/2,1/2}^o$	Ne II, Na III, Mg IV, Ar IV
$\dots np^6$	1S_0	Ne I, Na II, Mg III, Ar III


Cl, OI don't exist in HII regions (carbon is ionized)

NeV, MgV is too highly ionized

NII, OIII and SIII are useful temperature diagnostics

(Ne III and Ar III useful as well, but req higher energy photons)


Temperature Sensitive Line Ratios



at typical HII region densities,
NII transitions from ¹S₀ and ¹D₂
are below critical density

means:
approximately every collision
results in a radiative decay
(i.e. A wins over k)


Temperature Sensitive Line Ratios



Emission from 4-3 transition per NII =
 (rate of collisions that populate 4) x
 (fraction of radiative transitions in 4-3) x
 (energy of 4-3 transition)

$$P(4 \rightarrow 3) = \frac{n_e k_{04} \times A_{43}}{A_{43} + A_{41}} \times E_{43}$$


Temperature Sensitive Line Ratios



Emission from 3-2 transition per NII =
 (rate of collisions & radiative transitions
 that populate 3) x
 (fraction of radiative transitions in 3-2) x
 (energy of 3-2 transition)

$$P(3 \rightarrow 2) = \frac{n_e (k_{03} + k_{04} A_{43} / (A_{43} + A_{41})) \times A_{32} / (A_{32} + A_{31}) \times E_{32}}{A_{32} / (A_{32} + A_{31}) \times E_{32}}$$

Temperature Sensitive Line Ratios



Line Ratio:

$$\frac{P(4 \rightarrow 3)}{P(3 \rightarrow 2)} = \frac{A_{43}E_{43}}{A_{32}E_{32}} \left[\frac{k_{04}(A_{32} + A_{31})}{k_{03}(A_{43} + A_{41}) + k_{04}A_{43}} \right]$$

Define "collision strength" Ω_{ul}

$$k_{ul} = \frac{h^2}{(2\pi m_e)^{3/2}} \frac{1}{(kT)^{3/2}} \frac{\Omega_{ul}}{g_u}$$

separates gas temperature from
atomic properties

Detailed balance lets us get k_{ul} from k_{lu}

Temperature Sensitive Line Ratios

$$\frac{P(4 \rightarrow 3)}{P(3 \rightarrow 2)} = \frac{A_{43}E_{43}}{A_{32}E_{32}} \left[\frac{\Omega_{40}(A_{32} + A_{31})}{\Omega_{30}(A_{43} + A_{41})e^{E_{43}/kT} + \Omega_{40}A_{43}} \right]$$

Line ratio doesn't depend on density,
only on temperature.

Only density insensitive below the critical density.

Temperature Sensitive Line Ratios

